



## JMJD2B rabbit pAb

**Cat#: orb768169 (Manual)** 

For research use only. Not intended for diagnostic use.

Product Name JMJD2B rabbit pAb

Host species Rabbit

Applications IHC;IF;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

**Recommended dilutions** Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human KDM4B. AA range:351-400

Specificity JMJD2B Polyclonal Antibody detects endogenous levels of JMJD2B protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Lysine-specific demethylase 4B

Gene Name KDM4B

Cellular localization Nucleus.

**Purification** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Clonality** Polyclonal





Explore. Bioreagents.

1 mg/mlConcentration

**Observed band** 

**Human Gene ID** 23030

**Human Swiss-Prot Number** O94953


**Alternative Names** KDM4B; JHDM3B; JMJD2B; KIAA0876; Lysine-specific demethylase 4B;

JmjC domain-containing histone demethylation protein 3B; Jumonji domain-

containing protein 2B

cofactor:Binds 1 Fe(2+) ion per subunit.,domain:The 2 Tudor domains recognize and bind methylated histones. Double Tudor domain has an **Background** 

interdigitated structure and the unusual fold is required for its ability to bind methylated histone tails, function: Histone demethylase that specifically methylated histone tails., function: Histone demethylase that specifically demethylates 'Lys-9' of histone H3, thereby playing a role in histone code. Does not demethylate histone H3 'Lys-4', H3 'Lys-27', H3 'Lys-36' nor H4 'Lys-20'. Only able to demethylate trimethylated H3 'Lys-9', with a weaker activity than KDM4A, KDM4C and KDM4D. Demethylation of Lys residue generates formaldehyde and succinate., similarity: Belongs to the JHDM3 histone demethylase family., similarity: Contains 1 JmjC domain., similarity: Contains 1 JmjN domain., similarity: Contains 2 PHD-type zinc fingers., similarity: Contains 2 Tudor domains.,



Immunohistochemistry analysis of paraffin-embedded human brain, using JHD3B Antibody. The picture on the right is blocked with the synthesized